

**Case presentation**

**Mycophenolate mofetil therapy for pediatric bullous pemphigoid**

**Lucia Seminario-Vidal MD PhD<sup>1</sup>, Naveed Sami MD<sup>1</sup>, Jonathan Miller BS<sup>2</sup>, and Amy Theos MD<sup>1</sup>**

**Dermatology Online Journal 21 (8): 6**

**<sup>1</sup>Department of Dermatology, University of Alabama at Birmingham, Birmingham, AL**

**<sup>2</sup>Quillen College of Medicine, East Tennessee State University, Johnson City, TN**

**Corresponding author:**

Amy Theos, MD  
Children's South Specialty Clinic  
1940 Elmer J. Bissell Road, Birmingham, AL, 35243  
205-824-4823 (phone)  
205-824-4994 (fax)  
amy.theos@chsys.org (email)

---

**Abstract**


Bullous pemphigoid (BP) is a common autoimmune blistering disease in the adult population, but extremely rare in the pediatric population. Childhood BP usually has a favorable prognosis and responds well to topical and oral steroids. However, for patients that do not respond to corticosteroids, therapeutic alternatives are scarce. We report a case of a toddler with recalcitrant BP who was successfully treated with mycophenolate mofetil (MMF).

**Keywords: vesiculobullous diseases, bullous pemphigoid, mycophenolate mofetil, immunosuppressive therapy**

**Case synopsis**

A 16-month-old boy presented with worsening pruritic urticarial plaques and tense bullae (Figure 1) that began approximately one week after receiving his scheduled immunizations. Biopsies revealed a subepidermal bullous dermatosis with eosinophils (Figure 2), and direct immunofluorescence (DIF) was positive for strong linear IgG and C3 at the dermal epidermal junction. There was no significant positivity for IgM, IgA, or fibrin with appropriate reactive controls. Immunostaining for type IV collagen highlighted the basement membrane at the floor of the blister. Indirect immunofluorescence (IIF) was positive for linear deposits of IgG anti-basement membrane zone autoantibodies at the blister roof on salt-split skin. Staining on rodent bladder epithelium was negative. IIF did not reveal circulating IgA antibodies, excluding the possibility of linear IgA bullous dermatosis (LABD). Enzyme-linked immunosorbent assay (ELISA) for anti-BP180 autoantibody titers were elevated at 230 IU/ml. Other pertinent laboratory tests included a white blood cell count of 14.52 cells/mL (normal 5.98-13.51 cells/ml) and prominent eosinophilia of 3.13 cells/ml (normal 0.02-0.82 cells/ml). These clinical and laboratory findings were consistent with a diagnosis of BP. Thus, the patient was started on oral prednisone 1 mg/kg/day and topical steroids. Despite this treatment, his cutaneous disease worsened and he developed oral involvement, which led to an increase of the prednisone dose to 2 mg/kg/day and the addition of dapsone

with partial improvement. The patient was hospitalized for superinfection with *Staphylococcus aureus*. The infection was treated successfully with clindamycin and dapsone was discontinued. Following recovery of the acute infection, mycophenolate mofetil (MMF) was initiated at 30mg/kg/day. Because MMF needs several weeks to induce therapeutically relevant immunosuppression, prednisone was slowly tapered at a rate of 1mg/week over three months. Within two months of initiating this therapy his skin lesions cleared and the anti-BP180 antibodies decreased to 59 IU/ml. He received MMF for six months and, subsequently, it was slowly tapered down over 3 months; his skin remained clear (Figure 3). He did not experience any flares during MMF therapy or in the subsequent 4 months.



**Figure 1.** Tense bullae involving trunk and extremities. **Figure 2.** Subepidermal blister with eosinophils. (hematoxylin-eosin, original magnification x200)

## Discussion

Bullous pemphigoid (BP), initially described by Lever in 1953, is the most common autoimmune blistering disorder [1]. It is characterized immunologically by the presence of autoantibodies directed against 2 antigens found in the hemidesmosomes of the dermal-epidermal junction, i.e., BP antigen 1 (BPAG1 or AgBP230), and BPAG2 (or AgBP180 or collagen XVII) [2, 3]. BP primarily affects the elderly, but it may rarely occur in children as young as 2 months of age [4].

BP often starts as pruritus with urticarial or erythematous plaques that evolve over weeks to months into large, tense, or sometimes hemorrhagic bullae. Acral involvement is frequently seen in affected infants, whereas genital and oral involvement is more common in older children [5]. Ocular involvement may not occur until years after the onset of the condition and presents as dryness of the eyes with a feeling of chronic, intractable conjunctival irritation [5].

As with most autoimmune blistering diseases, the precise reason for autoantibody formation is unknown, although several HLA subclasses have been associated with an increased susceptibility to this disorder [6]. Drug intake and viral infections have been reported as possible triggering events [7]. Furthermore, it has been noted that BP in infancy may present shortly after vaccination [8, 9, 10]. However, this relationship is controversial and may be purely coincidental because the frequency of vaccination during childhood is high.

The differential diagnosis varies according to the stage of the disease. The initial pruritic, urticarial plaques need to be differentiated from atopic dermatitis, urticarial, or erythema multiforme. Bullous lesions can also be seen in impetigo, LABD, dermatitis herpetiformis, bullous



**Figure 3.** Residual hypopigmentation on areas of previous bullae.

systemic lupus erythematosus, and epidermolysis bullosa acquisita. Diagnosis is established by clinical, histological, and immunofluorescence analysis. Biopsies show subepidermal blisters with an eosinophilic infiltrate. DIF of perilesional skin samples shows linear deposition of mainly IgG or C3 along the basement membrane. Additional studies include IIF and ELISA. IIF reveals circulating IgG autoantibodies directed against BP180/230 glycoproteins binding linearly at the dermal-epidermal junction of monkey esophagus, and/or binding to the roof of an artificial blister cavity of salt-split human skin. ELISA detects IgG autoantibodies directed against the NC16A domain of BP180 [11].

BP treatment is based more on clinical experience than on controlled studies. Systemic corticosteroids are the mainstay of treatment. Sulfapyridine, dapsone, azathioprine, cyclosporine, and rituximab have been used as steroid-sparing agents and/or in corticosteroid-refractory BP [7, 12, 13]. Erythromycin with or without nicotinamide has been beneficial in some children, presumably owing to its anti-inflammatory effects [14]. In our patient, combined therapy of systemic corticosteroids and dapsone was unsuccessful.

Taking into account our patient's hospitalization for super-infection, his poor response to conventional treatment, and the need for therapy with minimal side effects, we decided to start mycophenolate mofetil (MMF). MMF is a prodrug of mycophenolic acid, which inhibits the rate-limiting enzyme in de novo synthesis of guanosine nucleotides, i.e., inosine monophosphate dehydrogenase. T and B lymphocytes are more dependent on this pathway than other cell types. Thus, MMF is effective at inhibiting T and B cell proliferation, inducing T cell apoptosis, and blocking antibody production by B cells [15]. This drug has previously been utilized in the treatment of adults with BP. It was found that MMF was equally effective to azathioprine in inducing disease remission (100%) when combined with corticosteroids. However, the MMF group had less liver toxicity [16]. In children, MMF therapy has also demonstrated to be safer when compared with other immunosuppressive agents for the treatment of atopic dermatitis, lupus nephritis, and nephritic syndrome [17-19]. This case illustrates that MMF could be a therapeutic alternative to traditional systemic immunosuppressive agents, with less favorable side-effect profiles, in treating refractory BP in children.

## References

1. Lever WF. Pemphigus. *Medicine (Baltimore)*. 1953 Feb;32(1):1-123. [PMID: 13024494]
2. Guidice GJ, Emery DJ, Diaz LA. Cloning and primary structural analysis of the bullous pemphigoid autoantigen BP180. *J Invest Dermatol*. 1992;99:243-50. [PMID: 1324962]
3. Tanaka T, Parry DA, Klaus-Kovtun V, et al. Comparison of molecularly cloned bullous pemphigoid antigen to desmoplakin I confirms that they define a new family of cell adhesion junction plaque proteins. *J Cell Biol*. 1991;266:12555-9. [PMID: 1712022]
4. Cunha PR, Thomazeski PG, Hipólito É, et al. Bullous pemphigoid in a 2-month-old infant. *Int J Dermatol*. 1998;37(12):935-938. [PMID: 9888337]
5. Waisbourd-Zinman O, Ben-Amitai D, Cohen AD, et al. Bullous pemphigoid in infancy: Clinical and epidemiologic characteristics. *J Am Acad Dermatol*. 2008 Jan;58(1):41-8. [PMID: 17945382]
6. Delgado JC, Turbay D, Yunis EJ, et al. A common major histocompatibility complex class II allele HLA-DQB1\*0301 is present in clinical variants of pemphigoid. *Proc Natl Acad Sci USA*. 1996;93:8569-8571. [PMID: 8710911]
7. Marcus KA, Halbertsma FJ, van Steensel MA. A case of juvenile bullous pemphigoid--successful treatment with diaminodiphenylsulfone and prednisone. *Pediatr Dermatol*. 2009 Jan-Feb;26(1):55-8. [PMID: 19250407]
8. Baykal C, Okan G, Sarica R. Childhood bullous pemphigoid developed after the first vaccination. *J Am Acad Dermatol*. 2001;44:348-350. [PMID: 11174412]
9. Erbagci Z. Childhood bullous pemphigoid following hepatitis B immunization. *J Dermatol*. 2002;29:781-785. [PMID: 12532044]
10. De la Fuente S, Hernández-Martín Á, de Lucas R, et al. Postvaccination bullous pemphigoid in infancy: report of three new cases and literature review. *Pediatr Dermatol*. 2013 Nov-Dec;30(6):741-4. [PMID: 24125034]
11. Sárdy M, Kostaki D, Varga R, et al. Comparative study of direct and indirect immunofluorescence and of bullous pemphigoid 180 and 230 enzyme-linked immunosorbent assays for diagnosis of bullous pemphigoid. *J Am Acad Dermatol*. 2013 Nov;69(5):748-53. [PMID: 23969034]
12. Thivolet J, Barthelemey H, Rigot-Muller G, et al. Effects of cyclosporin on bullous pemphigoid and pemphigus. *Lancet*. 1985;1:334-335. [PMID: 2857378]
13. Schulze J, Bader P, Henke U, et al. Severe bullous pemphigoid in an infant--successful treatment with rituximab. *Pediatr Dermatol*. 2008 Jul-Aug;25(4):462-5. [PMID: 18789089]
14. Oranje AP, Vuzevski VD, van Joost T, et al. Bullous pemphigoid in children: report of three cases with special emphasis on therapy. *Int J Dermatol*. 1991;30:339-42. [PMID: 1829723]
15. Eskin-Schwartz M, David M, Mimouni D. Mycophenolate mofetil for the management of autoimmune bullous diseases. *Dermatol Clin*. 2011 Oct;29(4):555-9. [PMID: 21924997]

16. Beissert S, Werfel T, Frieling U, et al. A comparison of oral methylprednisolone plus azathioprine or mycophenolate mofetil for the treatment of bullous pemphigoid. *Arch Dermatol.* 2007 Dec;143(12):1536-42. [PMID: 18087004]
17. Feng L, Deng J, Huo DM, et al. Mycophenolate mofetil versus azathioprine as maintenance therapy for lupus nephritis: a meta-analysis. *Nephrology (Carlton).* 2013 Feb;18(2):104-10. [PMID: 23113811]
18. Heller M, Shin HT, Orlow SJ, et al. Mycophenolate mofetil for severe childhood atopic dermatitis: experience in 14 patients. *Br J Dermatol.* 2007;157:127-32. [PMID: 17489974]
19. Gellermann J, Weber L, Pape L, et al. Gesellschaft für Pädiatrische Nephrologie (GPN). Mycophenolate mofetil versus cyclosporin A in children with frequently relapsing nephrotic syndrome. *J Am Soc Nephrol.* 2013 Oct;24(10):1689-97. [PMID: 23813218]