

# UC Davis

## Dermatology Online Journal

### Title

Preterm infant with a late presentation of blueberry muffin lesions secondary to recombinant erythropoietin

### Permalink

<https://escholarship.org/uc/item/4bm843q3>

### Journal

Dermatology Online Journal, 19(9)

### Authors

Rajpara, Anand  
Blackmon, Joseph  
Laarman, Rachel  
et al.

### Publication Date

2013

### DOI

10.5070/D3199019618

### Copyright Information

Copyright 2013 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

## Case Presentation

### Preterm infant with a late presentation of blueberry muffin lesions secondary to recombinant erythropoietin

Anand Rajpara MD<sup>1</sup>, Joseph Blackmon MD<sup>1</sup>, Rachel Laarman MD<sup>1</sup>, Robert Skaggs BS<sup>2</sup>, Ana Liolios MD<sup>1</sup>,  
Deede Liu MD<sup>1</sup>, Garth Fraga MD<sup>3</sup>

Dermatology Online Journal 19 (9): 9

<sup>1</sup>University of Kansas Medical Center, Department of Medicine, Division of Dermatology, Kansas City, KS

<sup>2</sup>University of Kentucky College of Medicine, Lexington, KY

<sup>3</sup>University of Kansas Medical Center, Department of Pathology and Laboratory Medicine, Kansas City, KS

## Correspondence:

Joseph Blackmon, MD  
PGY3 Dermatology Resident  
Department of Dermatology  
3901 Rainbow Boulevard  
Kansas City, Kansas 66160  
Tel: (913) 588-5000  
Email: jblackmon@kumc.edu

## Abstract

Our patient is a 26-week-old preterm female infant delivered by caesarean section secondary to severe maternal preeclampsia who had been receiving subcutaneous recombinant erythropoietin (r-EPO) for anemia of prematurity. At 8 weeks of age after 8 doses of r-EPO, the infant developed numerous non-blanching erythematous macules and patches located on the back, posterior shoulder, and posterior arms, concerning for late-onset blueberry muffin lesions. Biopsy of the lesions confirmed dermal hematopoiesis. After r-EPO was discontinued all skin lesions gradually resolved over a period of 2 weeks and never recurred.

**Keywords: recombinant erythropoietin, extramedullary dermal hematopoiesis, blueberry muffin lesions, anemia of prematurity**

## Case synopsis

Our patient is a 26-week-old preterm female infant delivered by caesarean section secondary to severe maternal preeclampsia. In addition to prematurity, the early clinical course was complicated by severe respiratory distress, severe intrauterine growth retardation, grade three intraventricular hemorrhage, anemia, and thrombocytopenia. Over the next several weeks the female infant's development progressed very well without any medical setbacks. However, at 8 weeks of age the patient suddenly developed an alarming rash on her back for which a dermatology consultation was consulted.

Prior to the onset of the rash the patient had been receiving subcutaneous recombinant erythropoietin (r-EPO) injections to treat anemia of prematurity at a dose of 400 units/kg three times a week for 9 doses. It was after the 8th dose of r-EPO that the rash appeared. Physical exam revealed numerous non-blanching erythematous macules and patches located on the infant's back, posterior shoulder, and posterior arms concerning for blueberry muffin lesions (Figure 1). A 4mm punch biopsy demonstrated polychromatophilic erythroblasts within the hypodermis with no immature myeloid cells, dysplastic erythroblasts, or viral inclusions, confirming the diagnosis of dermal hematopoiesis (Figure 2).

The neonatal ICU team suspected that the dermal hematopoiesis was secondary to r-EPO because they recently had a similar male preterm infant with late-onset blueberry muffin lesions secondary to r-EPO administration. After r-EPO was discontinued in our patient, her skin lesions gradually resolved over a period of 2 weeks and never recurred. The patient is now 9 months of age and continues to do well.



Figure 1. non-blanching erythematous macules




Figure 2. Polychromatophilic erythroblasts within the hypodermis

## Discussion

Blueberry muffin lesions typically present at birth or appear in the first few days of life as non-blanching blue-purple macules or dome-shaped papules that favor the trunk, head, and neck [1,2]. These lesions usually clear 3-6 months after birth and are the result of extramedullary dermal hematopoiesis [2]. Cutaneous biopsy shows aggregates of normoblastic erythroid precursors in the dermis of affected skin [2]. These lesions are most commonly seen in neonates with congenital infections, but have also

occurred in neonates with hemolytic conditions like hereditary spherocytosis, twin-twin transfusions, and blood group incompatibility disorders such as RH isoimmunization. These findings are also seen in neoplastic diseases such as neuroblastoma, langerhans cell histiocytosis, and congenital leukemia [2,3]. For the classic presentation, the most important part of the work up for blueberry muffin lesions is to rule out TORCH infections, which include congenital rubella, toxoplasmosis, syphilis, enterovirus, parvovirus B19, and cytomegalovirus. These can usually be detected by serologies, including IgM, or by viral PCR [4]. Other labs such as complete blood count can be helpful for hematologic malignancies. Skin biopsy confirms the diagnosis of extramedullary hematopoiesis.

Late-onset blueberry muffin lesions and dermal hematopoiesis have only been reported twice in neonates, once in a case report of a congenital rubella syndrome and the other as a result r-EPO administration [1,5]. To our knowledge there is only one published report of rEPO-induced blueberry muffin lesions other than the case we present here [5]. Our patient was a female born at 26 weeks gestational age that developed non-blanching erythematous macules on the back at 8 weeks of age after her 8<sup>th</sup> dose of r-EPO. Similarly, the previously reported male preterm infant was born at 22 4/7-weeks gestational age and developed purple non-blanching papules on the back at approximately 10 weeks of age after his 4<sup>th</sup> dose of r-EPO.

In preterm infants, the hematocrit drops after birth owing to expected breakdown of red blood cells and frequent blood draws. Erythropoietin in neonates is found to be low, so r-EPO was initially used to increase the levels in neonates and stimulate native red cell production [6]. Recombinant erythropoietin is also thought to reduce the need for red blood cell transfusions. The evidence for the use of r-EPO is somewhat mixed, but it has become a common tool for treating preterm infants with anemia [7]. As the age of viability is pushed earlier, it is important for the consulting dermatologist to be aware of the dermatologic effects of r-EPO in the preterm population.

## References

1. Vozza A, Tolone C, et al. Late onset Blueberry Muffin Syndrome following congenital rubella. *J Eur Acad Dermatol Venereol*. 2003 Mar;17(2):204-5. [PMID:12705753]
2. Mehta V, Balachandran C, Lonikar V. Blueberry muffin baby: a pictorial differential diagnosis. *Dermatol Online J*. 2008 Feb 28;14(2):8. [PMID:18700111]
3. Pizarro A, Elorza D, et al. Neonatal dermal erythropoiesis associated with severe rhesus immunization: amelioration by high-dose intravenous immunoglobulin. *Br J Dermatol*. 1995 Aug;133(2):334-6. [PMID:7547415]
4. Paller A, Mancini A. *Hurwitz Clinical Pediatric Dermatology*. 4<sup>th</sup> edition. Elsevier, 2011. p28-36.
5. Pandey V, Dummula K, Fraga G, Parimi P. Late-onset blueberry muffin lesions following recombinant erythropoietin administration in a premature infant. *J Pediatr Hematol Oncol*. 2012 Oct;34(7):534-5. [PMID:23007340]
6. Halvorsen S, Bechensteen AG. Physiology of erythropoietin during mammalian development. *Acta Paediatr Suppl*. 2002;91(438):17-26. [PMID:12477260]
7. Aher SM, Ohlsson A. Early versus late erythropoietin for preventing red blood cell transfusion in preterm and/or low birth weight infants. *Cochrane Database Syst Rev*. 2012 Oct 17. [PMID:23076909]