

Granuloma annulare-like eruption in chronic lymphocytic leukemia

Yasmin Hadian MS^{1,2}, Kourosh Beroukhim MD^{1,2}, Maxwell A Fung MD¹, Joshua M Schulman MD^{1,2}

Affiliations: ¹Department of Dermatology, School of Medicine, University of California-Davis, Sacramento, California, USA, ²Division of Dermatology, Veterans Administration Northern California Health Care System, Mather, California, USA

Corresponding Author: Joshua M. Schulman MD, Division of Dermatology, Building 801A, 10535 Hospital Way, Mather, CA 95655, Email: joshua.schulman@va.gov

Abstract

We report a patient with new-onset ulcerated granuloma annulare with concomitant involvement of B-cell leukemia. A granuloma annulare-like eruption with concomitant B cell chronic lymphocytic leukemia involvement in the skin is extremely rare, as only three cases have been previously reported in the literature to our knowledge. Given the rarity of ulceration in conventional granuloma annulare, it is possible this finding may serve as a diagnostic clue for underlying malignancy.

Keywords: granuloma annulare, granulomatous dermatitis, chronic lymphocytic leukemia, cutaneous leukemia, ulcer

Introduction

Granuloma annulare (GA) is a self-limited condition classically characterized by an asymptomatic eruption of annular, erythematous papules or plaques, most frequently affecting the distal extremities [1]. Ulceration is uncommon. Histopathological findings include interstitial or palisading histiocytes and lymphocytic infiltration in the dermis [1]. The pathogenesis has yet to be fully determined [2]. Although an association between GA and hematopoietic malignancies has been reported [3-13], GA-like infiltrates with concomitant involvement of B-cell chronic lymphocytic leukemia (B-CLL) in the same skin lesion is extremely rare [14,15]. We report a 75-year-old man who presented with an ulcerating GA-like eruption with cutaneous

B-CLL co-involvement in association with systemic B-CLL progression.

Case Synopsis

A 75-year-old man with Rai stage I-II B-CLL presented to the dermatology clinic for evaluation of a new skin eruption. He had a 10-year history of B-CLL, for which he was monitored without treatment. One month prior to presentation, he developed approximately five erythematous papules and plaques, several with central ulceration, on the back and abdomen that progressively enlarged. The patient reported "irritation" of the lesions upon friction against clothing, though they were otherwise asymptomatic. Around the time of onset, the patient had acute bronchitis that was treated with doxycycline. Review of systems was negative for fever, chills, and night sweats.

Physical examination revealed five discrete 0.5cm–3.0cm mildly edematous, erythematous papules and plaques on the mid-back (**Figure 1**), flanks, and anterior left forearm. The larger plaques demonstrated central ulceration with fibrinous crust (**Figure 1**). Clinical differential diagnoses included Sweet syndrome, cutaneous involvement of leukemia/lymphoma, and granulomatous dermatitis.

Histopathology revealed a mixed inflammatory infiltrate composed mostly of interstitial histiocytes, many multinucleated and exhibiting elastophagocytosis, and moderately dense perivascular/periadnexal lymphocytes (**Figure 2**),



Figure 1. Clinical findings. Physical examination revealed 0.5–3.0cm edematous, erythematous papule and plaque on the mid-back. The plaque demonstrated central ulceration with fibrinous crust.

the vast majority of which were CD20⁺ with variable CD5 co-expression.

A diagnosis of granuloma annulare-like dermatitis with concurrent cutaneous B-cell lymphocytic leukemia involvement was made. Upon 3-week follow-up, the lesions remained asymptomatic and the previously ulcerated lesions had re-epithelialized. The hematology-oncology department was consulted for evaluation and management of possible CLL progression. Within three months of initial presentation the patient developed systemic symptoms consisting of fever, night sweats, and fatigue, in association with progressively worsening lymphocytosis and cytopenia indicating systemic progression of his B-CLL.

Case Discussion

Although uncommon, cutaneous involvement by B-CLL may occur either as a primary leukemia cutis or as a secondary infiltrate associated with a separate neoplastic or inflammatory process [15-23]. The co-involvement of B-CLL with a GA-like infiltrate in the same cutaneous lesion is extremely rare [14,15]. To our knowledge, only three other cases have been reported (**Table 1**). One proposed mechanism for this unusual finding is that a granulomatous reaction may occur in response to locally infiltrating malignant lymphocytes [15]. Conversely, it is possible that neoplastic cells may be secondarily recruited to a site of primary dermatosis as a reactive inflammatory process [15]. For example, cutaneous B-CLL has been reported in healing varicella-zoster lesions [15,24,25]. In those cases, it is possible that Wolf isotopic response may have played a role, in which a specific cutaneous site that has been “immunocompromised” by a prior primary dermatosis may be subsequently affected by a secondary opportunistic pathology [25,26]. A similar phenomenon may be at play in our case.

In previously reported cases, patients with cutaneous co-involvement of granulomatous dermatitis and B-CLL presented with an eruption of erythematous, annular papules or plaques on the extremities or trunk [14,15]. In contrast, our patient




Figure 2. Histopathology findings revealed **A)** interstitial histiocytes accompanied by a moderately dense perivascular/periadnexal lymphocytic infiltrate. H&E, 40 \times . **B)** On higher power, there are multinucleated histiocytes that demonstrate elastophagocytosis (arrows). H&E, 200 \times .

Table 1. Reported cases of concomitant granuloma annulare-like infiltrate and cutaneous involvement of B-CLL in same skin lesion.

Patient [Ref]	Age/ Gender	B-CLL History	Location	Ulcerated	Elastophagocytosis	Immunophenotype of atypical lymphocytes	Treatment & Outcome
1 [14]	73/Male	10 years; Rai stage IV	Upper extremities	No	Yes	CD5, CD43, CD20, CD79a	Topical steroid, antihistamine, oral cephalexin, griseofulvin → no response; chemotherapy → cleared within 3 mo
2 [15]	72/Male	4 years; Rai stage I-II	Trunk, upper extremities	No	No	CD5, CD20, CD79a	TAC 0.1% cream, narrow band UV, and MTX sodium (no chemotherapy) → no response by 12 mo follow-up
3 [15]	76/Male	2 years; Rai stage I	Left arm	No	No	CD5, CD20, CD79a	None → did not clear by 24 mo follow-up
4 [present case]	75/Male	10 years; Rai stage I-II	Abdomen, flank, left upper extremity	Yes	Yes	CD5, CD20	None → improved but not cleared at 3 week follow-up

Ref, reference; B-CLL, B-cell chronic lymphocytic leukemia; TAC, triamcinolone acetonide; UV, ultraviolet; MTX, methotrexate.

uniquely demonstrated plaques that evolved with ulcerations. Histopathology typically reveals interstitial or palisading multinucleated histiocytes admixed with perivascular, atypical, small lymphocytes with small nucleoli throughout the dermis [14,15]. Immunohistochemistry may demonstrate atypical lymphocytes predominately expressing B cell markers CD20 and CD79a, as well as aberrant co-expression of T cell markers CD5 and CD43 [14,15].

Interestingly, histopathology of our patient's back lesions revealed elastophagocytosis, which is more commonly observed in actinic granuloma or annular elastolytic granuloma typically in sun-exposed areas [1,14,27-29]. However, the main sites of involvement in our patient were not sun-exposed.

The skin eruption in our patient was followed by progressive systemic signs of B-CLL, suggesting a possible correlation between this cutaneous manifestation and the systemic progression of B-CLL,

although this correlation has not been widely established. Only one of three other reported cases of concomitant B-CLL with a GA-like infiltrate also demonstrated systemic progression.

Interestingly, a previous case involving GA-like infiltrate with co-involvement of B-CLL reported clearance of the lesions after three months of chemotherapy [14], whereas cases of similar pathology that did not receive chemotherapy did not demonstrate resolution of the lesions [15]. Thus, chemotherapy may be an effective option for treating the GA-like lesions, in addition to the underlying malignancy, when presenting concomitantly with locally infiltrating malignant lymphocytes. Furthermore, these findings also indicate that a GA-like eruption in an elderly patient may suggest an underlying malignancy as a possible etiologic mechanism. Additional systemic diseases suggested to be associated with GA include diabetes mellitus, hyperlipidemia, sarcoidosis, Sweet

syndrome, thyroid disease, and infectious diseases including HIV/AIDS [2,30], highlighting the importance of a thorough clinical evaluation in patients with GA.

Conclusion

Granuloma annulare-like dermatitis with cutaneous B-CLL involvement is extremely rare and this is the first reported case with ulcerative morphology. Importantly, the eruption of GA-like lesions may

indicate underlying malignancy. Owing to the association of GA with systemic disease, it is essential for clinicians to further investigate a GA-like eruption, particularly in the elderly, with a thorough review of systems and workup. In patients with a known history of CLL, clinical suspicion should be appropriately elevated.

Potential conflicts of interest

The authors declare no conflicts of interest.

References

- Piette EW, and Rosenbach M. Granuloma annulare: clinical and histologic variants, epidemiology, and genetics. *J Am Acad Dermatol.* 2016;75:457-465. [PMID 27543209].
- Piette EW, and Rosenbach M. Granuloma annulare: pathogenesis, disease associations and triggers, and therapeutic options. *J Am Acad Dermatol.* 2016;75:467-79. [PMID 27543210].
- Barksdale SK, Perniciaro C, Halling KC, Strickler JG. Granuloma annulare in patients with malignant lymphoma: clinicopathologic study of thirteen new cases. *J Am Acad Dermatol.* 1994;31:42-48. [PMID 8021370].
- Wanat KA, Elenitsas R, Kim EJ, Rosenbach M. Granuloma annulare associated with cutaneous marginal zone lymphoma: a case linking a hematologic malignancy with granulomatous dermatitis. *Am J Dermatopathol.* 2012;34:844-846. [PMID 22935889].
- Dadban A, Slama B, Azzedine A, Lepeu G. Widespread granuloma annulare and hodgkin's disease. *Clin Exp Dermatol.* 2008;33:465-468. [PMID 18462441].
- Setoyama M, Kerdel FA, Byrnes JJ, Kanzaki T. Granuloma annulare associated with hodgkin's disease. *Int J Dermatol.* 1997;36:445-448. [PMID 9248891].
- Schwartz RA, Hansen RC, Lynch PJ. Hodgkin's disease and granuloma annulare. *Arch Dermatol.* 1981;117:185-186. [PMID 7212733].
- Bassi A, Scarfi F, Galeone M, Arunachalam M, Difonzo E. Generalized granuloma annulare and non-hodgkin's lymphoma. *Acta Derm Venereol.* 2013;93:484-485. [PMID 23250047].
- Martin JE, Wagner AJ, Murphy GF, Pinkus GS, Wang LC. Granuloma annulare heralding angioimmunoblastic T-cell lymphoma in a patient with a history of epstein-barr virus-associated B-cell lymphoma. *J Clin Oncol.* 2009;27:e168-e171. [PMID 19770367].
- Askin U, Durdu M, Senel E. Generalized granuloma annulare in a patient with myelocytic leukemia and chronic hepatitis B virus infection. *Indian J Dermatol Venereol Leprol.* 2009;75:287-289. [PMID 19439883].
- Hinckley MR, Walsh SN, Molnar I, Sheehan DJ, Sanguenza OP, Yosipovitch G. Generalized granuloma annulare as an initial manifestation of chronic myelomonocytic leukemia: a report of two cases. *Am J Dermatopathol.* 2008;30:274-277. [PMID 18496432].
- Cohen PR. Granuloma annulare, relapsing polychondritis, sarcoidosis, and systemic lupus erythematosus: conditions whose dermatologic manifestations may occur as hematologic malignancy-associated mucocutaneous paraneoplastic syndromes. *Int J Dermatol.* 2006;45:70-80. [PMID 16426383].
- Cohen PR. Granuloma annulare associated with malignancy. *South Med J.* 1997;90(10):1056-9. [PMID 9347823].
- Fullen DR, Jacobson SN, Valdez R, Novice FM, Lowe L. Granuloma annulare-like infiltrates with concomitant cutaneous involvement by B-cell non-hodgkin's lymphoma: report of a case. *Am J Dermatopathol.* 2003;25:57-61. [PMID 12544102].
- Sokumbi O, Gibson LE, Comfere NI, Peters MS. Granuloma annulare-like eruption associated with B-cell chronic lymphocytic leukemia. *J Cutan Pathol.* 2012;39:996-1003. [PMID 22830642].
- Plaza JA, Comfere NI, Gibson LE, et al. Unusual cutaneous manifestations of B-cell chronic lymphocytic leukemia. *J Am Acad Dermatol.* 2009;60:772-780. [PMID 19389520].
- Mehrany K, Byrd DR, Roenigk RK, et al. Lymphocytic infiltrates and subclinical epithelial tumor extension in patients with chronic leukemia and solid-organ transplantation. *Dermatol Surg.* 2003;29:129-34. [PMID 12562340].
- Dargent JL, Kornreich A, Andre L, Lespagnard L. Cutaneous infiltrate of chronic lymphocytic leukemia surrounding a primary squamous cell carcinoma of the skin. Report of an additional case and reflection on its pathogenesis. *J Cutan Pathol.* 1998;25:479-80. [PMID 9821078].
- Padgett JK, Parlette HL III, English JC III. A diagnosis of chronic lymphocytic leukemia prompted by cutaneous lymphocytic infiltrates present in mohs micrographic surgery frozen sections. *Dermatol Surg.* 2003;29:769-71. [PMID 12828704].
- Smoller BR, Warnke RA. Cutaneous infiltrate of chronic lymphocytic leukemia and its relationship to primary cutaneous epithelial neoplasms. *J Cutan Pathol.* 1998;25:160-4. [PMID 9550315].
- Wilson ML, Elston DM, Tyler WB, Marks VJ, Ferringer T. Dense lymphocytic infiltrates associated with non-melanoma skin cancer in patients with chronic lymphocytic leukemia. *Dermatol Online J.* 2010;16:4. [PMID 20233561].
- Agnew KL, Ruchlemer R, Catovsky D, Matutes E, Bunker CB. Cutaneous findings in chronic lymphocytic leukaemia. *Br J Dermatol.* 2004;150:1129-35. [PMID 15214899].
- Cerroni L, Zenahlik P, Hofler G, et al. Specific cutaneous infiltrates of B-cell chronic lymphocytic leukemia: a clinicopathologic and prognostic study of 42 patients. *Am J Surg Pathol.* 1996;20:1000-10. [PMID 8712287].
- Doutre MS, Beylot-Barry M, Beylot C, et al. Cutaneous localization

- of chronic lymphocytic leukemia at the site of chickenpox. *J Am Acad Dermatol.* 1997;36:98-9. [PMID 8996269].
25. Brasileiro A, Lencastre A, João A, Fidalgo A. Cutaneous leukemic infiltration following varicella - a case of wolf's isotopic response. *An Bras Dermatol.* 2016;91:72-75. [PMID 28300900].
 26. Piccolo V, Baroni A, Russo T, Schwartz RA. Ruocco's immunocompromised cutaneous district. *Int J Dermatol.* 2016;55:135-41. [PMID 26475059].
 27. Tock CL, Cohen PR. Annular elastolytic giant cell granuloma. *Cutis.* 1998;62:181-7. [PMID 9798106].
 28. Yaghoob R, Ranjbari N, Feily A. Actinic granuloma. *Dermatol Pract Concept.* 2014;4:31-32. [PMID 25126454].
 29. Chen WT, Hsiao PF, Wu YH. Spectrum and clinical variants of giant cell elastolytic granuloma. *Int J Dermatol.* 2017;56:738-745. [PMID 28239844].
 30. Cohen PR. Granuloma annulare: a mucocutaneous condition in human immunodeficiency virus-infected patients. *Arch Dermatol.* 1999;135:1404-7. [PMID 10566843].